Docket No.: PF-0475-2 DIV

APPENDIX

Claims on appeal:

- 11. An isolated antibody selected from the group consisting of:
- a) an antibody which specifically binds to a polypeptide comprising the amino acid sequence of SEQ ID NO:1, wherein the antibody specifically binds to an epitope of a polypeptide of SEQ ID NO:1,
- b) an antibody which specifically binds to a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to the amino acid sequence of SEQ ID NO:1, wherein said polypeptide has CoA dehydrogenase activity, and wherein the antibody specifically binds to an epitope of a polypeptide at least 90% identical to SEQ ID NO:1, and
- c) an antibody which specifically binds to a polypeptide comprising an immunogenic fragment of a polypeptide consisting of the amino acid sequence of SEQ ID NO:1, wherein said fragment comprises at least 15 contiguous amino acid residues of SEQ ID NO:1, and wherein the antibody specifically binds to an epitope of the fragment.
 - 31. The antibody of claim 11, wherein the antibody is:
 - a) a chimeric antibody,
 - b) a single chain antibody,
 - c) a Fab fragment,
 - d) a F(ab')₂ fragment, or
 - e) a humanized antibody.
 - 32. A composition comprising an antibody of claim 11 and an acceptable excipient.
 - 34. A composition of claim 32, wherein the antibody is labeled.

Docket No.: PF-0475-2 DIV

36. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 11, the method comprising:

- a) immunizing an animal with a polypeptide consisting of the amino acid sequence of SEQ ID NO:1, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
 - b) isolating antibodies from said animal, and
- c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide comprising the amino acid sequence of SEQ ID NO:1.
 - 37. A polyclonal antibody produced by a method of claim 36.
 - 38. A composition comprising the polyclonal antibody of claim 37 and a suitable carrier.
- 39. A method of making a monoclonal antibody with the specificity of the antibody of claim 11, the method comprising:
- a) immunizing an animal with a polypeptide consisting of the amino acid sequence of SEQ ID NO:1, or an immunogenic fragment thereof, under conditions to elicit an antibody response,
 - b) isolating antibody producing cells from the animal,
- c) fusing the antibody producing cells with immortalized cells to form monoclonal antibodyproducing hybridoma cells,
 - d) culturing the hybridoma cells, and
- e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide comprising the amino acid sequence of SEQ ID NO:1.
 - 40. A monoclonal antibody produced by a method of claim 39.
 - 41. A composition comprising the monoclonal antibody of claim 40 and a suitable carrier.

Docket No.: PF-0475-2 DIV

42. The antibody of claim 11, wherein the antibody is produced by screening a Fab expression library.

- 43. The antibody of claim 11, wherein the antibody is produced by screening a recombinant immunoglobulin library.
- 58. An isolated antibody which specifically binds to a polypeptide selected from the group consisting of:
 - a) a polypeptide consisting of the amino acid sequence of SEQ ID NO:1, and
- b) an immunogenic fragment of a polypeptide consisting of the amino acid sequence of SEQ ID NO:1, wherein said fragment comprises at least 15 contiguous amino acid residues of SEQ ID NO:1.